



**P 202** - Topic: Genetics and Genomics of Biotic and Abiotic Stress Resistance

## **Resistance of European winter wheat cultivars to *Zymoseptoria tritici* isolate IPO88004**

Paweł Czembor, Magdalena Radecka-Janusik

Plant Breeding & Acclimatization Institute, National Research Institute, Department of Plant Breeding and Genetics, Radzików, 05-870 Błonie, Poland

 Paweł Czembor     p.czembor@ihar.edu.pl

**Key message:** Resistance to *Septoria tritici* blotch in European cultivars is contributed mainly by quantitative loci and those with main effects.

*Septoria tritici* blotch (STB) of wheat (*Triticum aestivum*), caused by the fungal pathogen *Mycosphaerella graminicola* (anamorph: *Zymoseptoria tritici*, syn. *Septoria tritici*), is present in most wheat-growing areas worldwide. Host resistance is the most economical and safest method of controlling the disease and information on resistance loci is crucial for effective breeding for resistance programs. In the study we used a set of 83 wheat cultivars registered in the Descriptive List of Agricultural Plant Varieties (COBORU 2012), 92 cultivars from other European countries and 25 cultivars/lines with identified STB resistance loci. The wheat genotypes were tested on adult plant stage under polytunnel conditions with watering system. Fully expanded flag leaves were sprayed with spore suspension of IPO88004 *Z. tritici* isolate. After incubation period, the percentage leaf area covered by necrosis (NEC) and covered by pycnidia (PYC) were measured on flag leaf of each wheat cultivar/line that were used in agglomerative hierarchical clustering (AHC) analysis with UPGA algorithm (unweighted pair-group average). Six groups of wheat cultivars/lines were identified and the largest group comprised 132 resistant genotypes with average NEC 18.7% and PYC 8.0%. Within this group, set of 43 highly resistant wheat cultivars were identified (NEC min. 1.1% and max. 15.1%, PYC min. 0.4%; max. 8.1%): Tuareg, Salutos, Samurai, Florett, Intro, Capone, Grapeli, Carroll, Praktik, Tabasco, Chilton, RGT Kilimanjaro, Zappa, Belenus, Joker, Butaro, Kalahari, KWS Dacanto, Riband, Fregata, Elixer, Forum, Estivus, Meteor, Glaucus, Bockris, Satyna, RGT Djoko, Jenga, Heros, Olivin, Zobel, Magnus, Kredo, Kepler, Pamier, Pengar, Julius, Desamo, Gordian, Eron, Pionier, Lear, Mandub and Kranich. In addition in the same subgroup were classified three cultivars with identified resistance loci: Florett (*QTL-3B*, *QTL-6D*; *Stb6+Stb15*), Tuareg (*QTL-4B*, *QTL-6B*; *Stb6*) and Riband (*QTL-6B*). This may suggest that resistance to STB in European cultivars is contributed mainly by quantitative loci and those with main effects. Presented work (phenotyping data) is a part of larger project aiming at identification of resistance genes (*Stb*) to *Septoria tritici* blotch in winter wheat and will be used in near future in association mapping approach.

### Acknowledgments

This research was supported financially by the Polish Ministry of Agriculture and Rural Development, program Fundamental Research for Biological Progress in Crop Production (years 2014-2020).